


WHAT IS CLAIMED IS:

1. A compound represented by Formula (I):



(I)

or a pharmaceutically acceptable salt thereof, wherein:

X and Y each independently is aryl or heteroaryl wherein at least one of X and Y is a heteroaryl with N adjacent to the position of attachment to A or B respectively;

three of A¹, A², A³, A⁴, and A⁵ are N, the remaining are C, and one of A¹ and A⁴ must be N, but not both A¹ and A⁴ are N;

W is -C₃₋₇cycloalkyl, -heteroC₃₋₇cycloalkyl, -C₀₋₆alkylaryl, or -C₀₋₆alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO₂, -C₁₋₆alkyl, -C₁₋₆alkenyl, -C₁₋₆alkynyl, -OR¹, -NR¹R², -C(=NR¹)NR²R³, -N(=NR¹)NR²R³, -NR¹COR², -NR¹CO₂R², -NR¹SO₂R⁴, -NR¹CONR²R³, -SR⁴, -SOR⁴, -SO₂R⁴, -SO₂NR¹R², -COR¹, -CO₂R¹, -CONR¹R², -C(=NR¹)R², or -C(=NOR¹)R² substituents;

X is optionally substituted with 1-7 independent halogen, -CN, NO₂, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -OR¹, -NR¹R², -C(=NR¹)NR²R³, -N(=NR¹)NR²R³, -NR¹COR², -NR¹CO₂R², -NR¹SO₂R⁴, -NR¹CONR²R³, -SR⁴, -SOR⁴, -SO₂R⁴, -SO₂NR¹R², -COR¹, -CO₂R¹, -CONR¹R², -C(=NR¹)R², or -C(=NOR¹)R² substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C₁₋₆alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C₁₋₆alkyl, -O(C₀₋₆alkyl), -O(C₃₋₇cycloalkyl), -O(aryl), -O(heteroaryl), -N(C₀₋₆alkyl)(C₀₋₆alkyl), -N(C₀₋₆alkyl)(C₃₋₇cycloalkyl), or -N(C₀₋₆alkyl)(aryl) groups;

R¹, R², and R³ each independently is -C₀₋₆alkyl, -C₃₋₇cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C₁₋₆alkyl, -O(C₀₋₆alkyl), -O(C₃₋₇cycloalkyl), -O(aryl), -O(heteroaryl), -N(C₀₋₆alkyl)(C₀₋₆alkyl), -N(C₀₋₆alkyl)(C₃₋₇cycloalkyl), or -N(C₀₋₆alkyl)(aryl) substituents;

R⁴ is -C₁₋₆alkyl, -C₃₋₇cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C₁₋₆alkyl, -O(C₀₋₆alkyl), -O(C₃₋₇cycloalkyl), -O(aryl), -O(heteroaryl), -N(C₀₋₆alkyl)(C₀₋₆alkyl), -N(C₀₋₆alkyl)(C₃₋₇cycloalkyl), or -N(C₀₋₆alkyl)(aryl) substituents;

A is -C₀₋₄alkyl, -C₀₋₂alkyl-SO-C₀₋₂alkyl-, -C₀₋₂alkyl-SO₂-C₀₋₂alkyl-, -C₀₋₂alkyl-CO-C₀₋₂alkyl-, -C₀₋₂alkyl-NR⁹CO-C₀₋₂alkyl-, -C₀₋₂alkyl-NR⁹SO₂-C₀₋₂alkyl- or -heteroC₀₋₄alkyl;

Y is optionally substituted with 1-7 independent halogen, -CN, NO₂, -C₁₋₆alkyl, -C₂₋₆alkenyl, -C₂₋₆alkynyl, -OR⁵, -NR⁵R⁶, -C(=NR⁵)NR⁶R⁷, -N(=NR⁵)NR⁶R⁷, -NR⁵COR⁶, -NR⁵CO₂R⁶, -NR⁵SO₂R⁸, -NR⁵CONR⁶R⁷, -SR⁸, -SOR⁸, -SO₂R⁸, -SO₂NR⁵R⁶, -COR⁵, -CO₂R⁵, -CONR⁵R⁶, -C(=NR⁵)R⁶, or -C(=NOR⁵)R⁶ substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C₁₋₆alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C₁₋₆alkyl, -O(C₀₋₆alkyl), -O(C₃₋₇cycloalkyl), -O(aryl), -O(heteroaryl), -N(C₀₋₆alkyl)(C₀₋₆alkyl), -N(C₀₋₆alkyl)(C₃₋₇cycloalkyl), or -N(C₀₋₆alkyl)(aryl) groups;

R⁵, R⁶, and R⁷ each independently is -C₀₋₆alkyl, -C₃₋₇cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C₁₋₆alkyl, -O(C₀₋₆alkyl), -O(C₃₋₇cycloalkyl), -O(aryl), -O(heteroaryl), -N(C₀₋₆alkyl)(C₀₋₆alkyl), -N(C₀₋₆alkyl)(C₃₋₇cycloalkyl), or -N(C₀₋₆alkyl)(aryl) substituents;

R⁸ is -C₁₋₆alkyl, -C₃₋₇cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C₁₋₆alkyl, -O(C₀₋₆alkyl), -O(C₃₋₇cycloalkyl), -O(aryl), -O(heteroaryl), -N(C₀₋₆alkyl)(C₀₋₆alkyl), -N(C₀₋₆alkyl)(C₃₋₇cycloalkyl), or -N(C₀₋₆alkyl)(aryl) substituents;

B is -C₀₋₄alkyl, -C₀₋₂alkyl-SO-C₀₋₂alkyl-, -C₀₋₂alkyl-SO₂-C₀₋₂alkyl-, -C₀₋₂alkyl-CO-C₀₋₂alkyl-, -C₀₋₂alkyl-NR¹⁰CO-C₀₋₂alkyl-, -C₀₋₂alkyl-NR¹⁰SO₂-C₀₋₂alkyl-, or -heteroC₀₋₄alkyl;

R⁹ and R¹⁰ each independently is -C₀₋₆alkyl, -C₃₋₇cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C₁₋₆alkyl, -O(C₀₋₆alkyl), -O(C₃₋₇cycloalkyl), -O(aryl), -O(heteroaryl), -N(C₀₋₆alkyl)(C₀₋₆alkyl), -N(C₀₋₆alkyl)(C₃₋₇cycloalkyl), -N(C₀₋₆alkyl)(aryl) substituents;

Z is -C₃₋₇cycloalkyl, -heteroC₃₋₇cycloalkyl, -C₀₋₆alkylaryl, or -C₀₋₆alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO₂, -C₁₋₆alkyl, -C₁₋₆alkenyl, -C₁₋₆alkynyl, -OR¹, -NR¹R², -C(=NR¹)NR²R³, -N(=NR¹)NR²R³, -NR¹COR², -NR¹CO₂R², -NR¹SO₂R⁴, -NR¹CONR²R³, -SR⁴, -SOR⁴, -SO₂R⁴, -SO₂NR¹R², -COR¹, -CO₂R¹, -CONR¹R², -C(=NR¹)R², or -C(=NOR¹)R² substituents;

R¹¹ is halogen, -C₀₋₆alkyl, -C₀₋₆alkoxyl, =O, =N(C₀₋₄alkyl), or -N(C₀₋₄alkyl)(C₀₋₄alkyl);

any alkyl optionally substituted with 1-5 independent halogen substituents;
 any N may be an N-oxide;
 and one of W and Z is optionally absent.

5

2. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof,
 wherein:

X is 2-pyridyl optionally substituted with 1-4 independent halogen, -CN, NO₂, -C₁-
 6alkyl, -C₂-6alkenyl, -C₂-6alkynyl, -OR¹, -NR¹R², -C(=NR¹)NR²R³, -N(=NR¹)NR²R³,
 10 -NR¹COR², -NR¹CO₂R², -NR¹SO₂R⁴, -NR¹CONR²R³, -SR⁴, -SOR⁴, -SO₂R⁴, -SO₂NR¹R², -COR¹,
 -CO₂R¹, -CONR¹R², -C(=NR¹)R², or -C(=NOR¹)R² substituents, wherein optionally two substituents
 are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C₁-6alkyl
 substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5
 independent halogen, -CN, -C₁-6alkyl, -O(C₀-6alkyl), -O(C₃-7cycloalkyl), -O(aryl), -O(heteroaryl), -
 15 N(C₀-6alkyl)(C₀-6alkyl), -N(C₀-6alkyl)(C₃-7cycloalkyl), or -N(C₀-6alkyl)(aryl) groups.

3. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof,
 wherein:

Y is phenyl optionally substituted with 1-5 independent halogen, -CN, NO₂, -C₁-6alkyl,
 20 -C₂-6alkenyl, -C₂-6alkynyl, -OR⁵, -NR⁵R⁶, -C(=NR⁵)NR⁶R⁷, -N(=NR⁵)NR⁶R⁷, -NR⁵COR⁶,
 -NR⁵CO₂R⁶, -NR⁵SO₂R⁸, -NR⁵CONR⁶R⁷, -SR⁸, -SOR⁸, -SO₂R⁸, -SO₂NR⁵R⁶, -COR⁵, -CO₂R⁵, -
 CONR⁵R⁶, -C(=NR⁵)R⁶, or -C(=NOR⁵)R⁶ substituents, wherein optionally two substituents are
 combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C₁-6alkyl substituent,
 cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent
 25 halogen, -CN, -C₁-6alkyl, -O(C₀-6alkyl), -O(C₃-7cycloalkyl), -O(aryl), -O(heteroaryl), -N(C₀-
 6alkyl)(C₀-6alkyl), -N(C₀-6alkyl)(C₃-7cycloalkyl), or -N(C₀-6alkyl)(aryl) groups.

4. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof,
 wherein:

30 Z is -C₀-6alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN,
 NO₂, -C₁-6alkyl, -C₁-6alkenyl, -C₁-6alkynyl, -OR¹, -NR¹R², -C(=NR¹)NR²R³, -N(=NR¹)NR²R³, -
 NR¹COR², -NR¹CO₂R², -NR¹SO₂R⁴, -NR¹CONR²R³, -SR⁴, -SOR⁴, -SO₂R⁴, -SO₂NR¹R², -COR¹,
 -CO₂R¹, -CONR¹R², -C(=NR¹)R², or -C(=NOR¹)R² substituents; R¹¹ is halogen, -C₀-6alkyl, -C₀-
 6alkoxyl, =O, =N(C₀-4alkyl), or -N(C₀-4alkyl)(C₀-4alkyl).;

35

5. The compound according to Claim 1, consisting of

- 2-[4-(3-Methoxy-4-pyridin-2-ylphenyl)-2H-1,2,3-triazol-2-yl]pyridine;
2-[4-(3-methoxy-4-pyridin-2-ylphenyl)-1H-1,2,3-triazol-1-yl]pyridine;
2-[4-(3-pyridin-2-ylphenyl)-1H-1,2,3-triazol-1-yl]pyridine;
2-[4-(3-pyridin-2-ylphenyl)-2H-1,2,3-triazol-2-yl]pyridine;
5 2-[4-(3-pyridin-3-ylphenyl)-1H-1,2,3-triazol-1-yl]pyridine;
2-[4-(3-pyridin-3-ylphenyl)-2H-1,2,3-triazol-2-yl]pyridine;
2-[4-(3-fluoro-4-pyridin-2-ylphenyl)-1H-1,2,3-triazol-1-yl]pyridine;
2-[4-(3-fluoro-4-pyridin-2-ylphenyl)-2H-1,2,3-triazol-2-yl]pyridine;
2-[2-methoxy-4-(5-methyl-1-pyridin-2-yl-1H-1,2,3-triazol-4-yl)phenyl]pyridine;
10 2-[2-methoxy-4-(5-methyl-2-pyridin-2-yl-2H-1,2,3-triazol-4-yl)phenyl]pyridine.

15

or a pharmaceutically acceptable salt thereof.

20

6. A pharmaceutical composition comprising: a therapeutically effective amount of the compound according to claim 1, or a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable carrier.

25

7. The pharmaceutical composition according to claim 6, further comprising i) an opiate agonist, ii) an opiate antagonist, iii) a calcium channel antagonist, iv) a 5HT receptor agonist, v) a 5HT receptor antagonist, vi) a sodium channel antagonist, vii) an NMDA receptor agonist, viii) an NMDA receptor antagonist, ix) a COX-2 selective inhibitor, x) an NK1 antagonist, xi) a non-steroidal anti-inflammatory drug, xii) a GABA-A receptor modulator, xiii) a dopamine agonist, xiv) a dopamine
30 antagonist, xv) a selective serotonin reuptake inhibitor, xvi) a tricyclic antidepressant drug, xvii) a norepinephrine modulator, xviii) L-DOPA, xix) buspirone, xx) a lithium salt, xxi) valproate, xxii) neurontin, xxiii) olanzapine, xxiv) a nicotinic agonist, xxv) a nicotinic antagonist, xxvi) a muscarinic agonist, xxvii) a muscarinic antagonist, xxviii) a selective serotonin and norepinephrine reuptake inhibitor (SSNRI), xxix) a heroin substituting drug, xxx) disulfiram, or xxxi) acamprosate.

35

8. The pharmaceutical composition according to claim 7, wherein said heroin substituting drug is methadone, levo-alpha-acetylmethadol, buprenorphine or naltrexone.

9. The use of the compound of Claim 1 for the preparation of a medicament useful in the treatment of pain disorders, extrapyramidal motor function disorders, anxiety disorders, Parkinson's disease, depression, epilepsy, cognitive dysfunction, drug addiction, circadian rhythm and sleep disorders, and obesity.

10. The use according to claim 9 wherein said pain disorder is acute pain, persistent pain, chronic pain, inflammatory pain, or neuropathic pain.

11. The use of the compound of Claim 1 for the preparation of a medicament useful in the treatment of anxiety, depression, bipolar disorder, psychosis, drug withdrawal, tobacco withdrawal, memory loss, cognitive impairment, dementia, Alzheimer's disease, schizophrenia or panic.

12. The use according to claim 9 wherein said disorder of extrapyramidal motor function is Parkinson's disease, progressive supramuscular palsy, Huntington's disease, Gilles de la Tourette syndrome, or tardive dyskinesia.